



**West African Ornithological Society
Société d'Ornithologie de l'Ouest
Africain**



**Join the WAOS and support
the future availability of free
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.

If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la
SOOA et soutenez la
disponibilité future des pdfs
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.

Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante
(2) pour économiser du papier.

British Museum (Nat. Hist.) as Apaloderma narina brachyurum. I have subsequently seen this bird near Azare, and Philip Blasdale has reported it near Potiskum.

P Blasdale also reported Pygmy Flycatchers Alseonax minimus as being present in the Ruriwai area just off the Jos/Zaria road. Unfortunately no skins were procured to establish the race.

I have records of an additional species which can positively be added to the Plateau list. On 6th May 1952 I saw an African Blue Quail Excalfactoria adansoni near Sabon Gida Dan, and on 7th June 1960 I saw three birds of this species in practically the same place.

R.E. Sharland.

Red-necked Kestrel Falco chiquera hunting bats. For eight successive evenings until the time of writing (18th January 1965) I and others have watched a pair of Red-necked Kestrels taking bats shortly after their emergence from houses on the Ahmadu Bello University campus, Zaria. On each one of these evenings, one or both birds have been seen either striking at or carrying in the talons a bat, and I have seen the actual mid-air strike three or four times now.

The bats are of one or two fairly small species, up to 10" or 12" wingspread, and emerge from the eves en masse about 15 minutes after the sun sets. At about the time of emergence, or a few minutes before, i.e. about 1830 hrs., the falcons appear and sit quietly waiting on one of two or three favourite tree-top perches. No sooner do the bats emerge than the falcons set off in pursuit in fast powerful flight usually about tree-top level, but rising if necessary to one or two hundred feet. Just before the falcon closes on it, the bat takes avoiding action and as often as not escapes; although I had the impression that the birds sometimes do not try too hard. Once a bat was caught, held in the talons for a few seconds and then released. Having captured its prey, the kestrel makes off and doubtless perches to consume it. The performance is over by 1840 hrs.

Cave & MacDonald (Birds of the Sudan) say that the Red-necked Kestrel is a "courageous little falcon usually seen dashing after prey at breakneck speed." In the best tradition of the genus Falco, it certainly is a bold and fast flier, and habitually takes prey on the wing - weaver-birds, swifts and doves (Bannerman, 1954), and also grasshoppers, locusts and termites (Roberts, The Birds of South Africa, who also records rats and lizards in the diet). Mackworth-Praed & Grant (African Handbook of Birds) say that this kestrel is active at dusk, and I would venture to suggest that the falcon which Macgregor reported seeing catching bats at Kaduna (Bull. N.O.S. 3: 15) was not a Hobby F. subbuteo but the present species F. chiquera.

C.H. Fry