



**West African Ornithological Society
Société d'Ornithologie de l'Ouest
Africain**



**Join the WAOS and support
the future availability of free
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.
If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la
SOOA et soutenez la
disponibilité future des pdfs
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.
Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante
(2) pour économiser du papier.

in Sahel Savannah. On 28th June a small flock of White-throated Bee-eaters was seen on the River Ngadda 4 miles to the East of Maiduguri, and I am sure these were non-breeders.

Credit for the first Nigerian nest goes to Mrs J. Hopson who found one attended by adults in a garden on the South bank of the River Yobe - actually a near-stagnant stream or ditch at this time of year - at Yo about the end of June. During the subsequent fortnight, I found a further 15 or so nests in the area, mainly at Malam'fatori on Lake Chad, with some near the motor track between here and Damarsac. Breeding was in full swing, with some clutches on the point of hatching or in other cases up to six (the commonest number) moderately or well grown nestlings. Nests were extremely easy to find, since the country is very open and adults, especially when feeding nestlings, can be very easily watched to the nest. All nests were solitary, in flat or gently sloping sandy soil. An intensive study was made during the short time available and details of further findings will be published elsewhere. C.H. Fry

Grey-backed Fiscal Shrike in Nigeria. On 3rd July 1965 I found a pair (male and female) of Grey-backed Fiscal Shrikes Lanius excubitorius inhabiting scattered trees between quite open fallow fields and a swamp with tall grasses and dead-looking 20- and 40-ft. trees, a mile behind the shore of Lake Chad at Malam'fatori, North-East Bornu Province. Mr and Mrs A.J. Hopson and I saw the birds in exactly the same place again on 8th July. A moderately deep cup nest, 7 inches in diameter, constructed of small thorny twigs and some bents, and securely sited at 8 feet in a 10 ft. thorn shrub, reminded me of the nest of Lanius collurio and may well have belonged to the present species. The nest had been occupied this year, and both adult birds perched in the thorn bush. No 'larder' was found. Bannerman (1953) does not record it West of Lake Chad, but it is included in the Provisional Check-list of the birds of Nigeria (Elgood, 1964, Bull. N.O.S. 1 : 13-25. Gray informs me that he recorded this species, without taking any field notes, on the Mora - Waza road, Northern Cameroun, in February 1962. C.H. Fry

Guinea Wood-Hoopoe on the Jos Plateau. A Guinea Wood-Hoopoe Phoeniculus erythrophynchus which I recorded at Gindin, near Panyam, in 1959 is an addition to the provisional check-list of the birds of Plateau Province over 2,000 ft. (Ebbutt et al., 1964, Bull. N.O.S. 3 : 9-14), J. Lang

New records for Obudu Plateau. The following birds are not mentioned by Elgood (1965 : The Birds of the Obudu Plateau, Eastern Region of Nigeria; Nig. Field 30 : 60-69) but were recorded there by myself on two recent visits : Black Kite Milvus migrans (2 over grassland on 17th December, 1964); Leaf-Love Pyrhurus scandens (one on the forest edge, 18th December 1964); Scarlet-spectacled Wattle-eye Platysteira cyanea, White-crowned Cliff-Chat Thamnotaea coronata (on approach road at 4,200 ft.), and Pin-tailed Whydah Vidua macroura all in August 1965.

H.H. Gray