



**West African Ornithological Society  
Société d'Ornithologie de l'Ouest  
Africain**



**Join the WAOS and support  
the future availability of free  
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.

If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la  
SOOA et soutenez la  
disponibilité future des pdfs  
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.

Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante  
(2) pour économiser du papier.

Sooty Tern Recovery. An immature female Sooty Tern Sterna fuscata marked U.S. Fish & Wildlife Service 873-58863 was killed by a student of Molusi College, Ijebu-Igbo ( $6^{\circ} 58' N$ ,  $4^{\circ} 00' E$ ) some time between mid-April and late May, 1966. The skin is in my possession, but marking and recovery details have not yet been established.

D.R.Wells

Passage Dates of the White-throated Bee-eater at Kaduna. The table shows the first and last dates for northward and southward passage of White-throated Bee-eaters Aerops albicollis at Kaduna in five years. The regularity of this migration is remarkable.

Year	Northward passage		Southward passage	
	First	Last	First	Last
1962	6th May	23rd May	8th Oct	14th Nov
1963	11th May	24th May	9th Oct	13th Nov
1964	20th April	21st May	10th Oct	13th Nov
1965	4th May	25th May	10th Oct	12th Nov
1966	(Not observed)		27th Oct	22nd Nov*

\* Rains ended late.

Flights were all strictly North/South, except when seeking evening roosts; they were mainly in the morning and evening, with a marked decrease at mid-day. Most sightings refer to one to three flocks of 10-40 birds each; such multiple flocks usually have a common East/West front, and flocks are separated by 20 to 60 yards. They fly at approximately 30 to 200 ft. high, at an estimated 15/20 M.p.h. Purposeful migration is interrupted (a) by random circling for periods up to 2 minutes, and (b) by descent to tree tops for sporadic feeding of 20 seconds to 20 minutes duration. There have been periods of up to 4 days when no birds have been seen, but there is no evidence of specific waves (cf. Fry, 1964, Bull. Niger. Orn. Soc. 1 (3) : 6).

M.Skilleter

Status of the Bluethroat in Nigeria. The only Nigerian records of the Bluethroat Cyanosylvia svecica up to 1965 are given by Elgood et al. (1966, Ibis 108 : 84-116) i.e. one at Sokoto, 27th December 1953, one at Kano 23rd January 1955, one at Potiskum February 1959, and two Zaria, 3rd December 1962. The authors note that the species has recently been found to be quite common in Senegal, but conclude that "it seems unlikely that the Bluethroat is as common a visitor to Nigeria as it appears to be to Senegal".