



**West African Ornithological Society**  
**Société d'Ornithologie de l'Ouest**  
**Africain**



**Join the WAOS and support  
the future availability of free  
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.

If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la  
SOOA et soutenez la  
disponibilité future des pdfs  
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.

Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante  
(2) pour économiser du papier.

OLIVE-TREE WARBLER Hippolias olivetorum

AT KANO: A SPECIES NEW TO WEST AFRICA.

by R.E.Sharland.

The harmattan started early in Nigeria (17th. October 1971) and easterly winds prevailed for the following week. On 22nd. October I caught a striking warbler in my nets which on examination proved to be an Olive-tree Warbler Hippolias olivetorum. The upper parts were grey with a small white supercilium and the under parts white. The head had a very marked Hippolias-type peak and the bill was long and flattened. There was a very distinct wing panel caused by the pale outer webs of the secondaries. The outer tail feathers had pale tips. Measurements were: Wing: 83mm. Tail: 62mm. Bill: 18mm. Tarsus: 23mm. The wing formula was as follows: Wing point (longest primary) 3rd., 4th -1mm., 5th -4mm., 2nd -5mm. The bird was ringed and released. This appears to be the first record of the Olive-tree Warbler in West Africa.

.....

FURTHER OBSERVATIONS OF BUFF-HEADED

WOOD HOOPOE Phoeniculus bollei.

by Netta Robinson.

R.A.Honeywell (Some Interesting Observations from Ghana June 1971. BNOS 8:31) remarks on the appearance of Phoeniculus bollei in Ashanti.

I have records from Ilesha in Western Nigeria in 1964 of the same species in which the buff on the face, forehead and throat was distributed in the manner described by Honeywell (op.cit.). The birds which I observed, quite closely and frequently over a period of one year, also had red bills, I did not see any with horn coloured bills, and they moved about in groups of three. I have no record of the colour of the feet.

At the time, after consideration, I assumed them to be P.b.okuensis although, as pointed out by Honeywell, Bannerman (1953, Birds of West & Equatorial Africa. Vol. I.) regards this race as being restricted to the montane forest area of Lake Oku, in Cameroun.