

### West African Ornithological Society Société d'Ornithologie de l'Ouest Africain



## Join the WAOS and support the future availability of free pdfs on this website.

### http://malimbus.free.fr/member.htm

If this link does not work, please copy it to your browser and try again. If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

# Devenez membre de la SOOA et soutenez la disponibilité future des pdfs gratuits sur ce site.

#### http://malimbus.free.fr/adhesion.htm

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer. Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante (2) pour économiser du papier.

Specimens in the Zoology Museum, Univ. Ibadan, from Gambari Forest Reserve, Western Nigeria, are typical nominate race P.b.bollei. It should also be remembered that okuensis has no buff colouration on the sides of the face. Ed.

### LONG-LEGGED BUZZARDS Buteo rufinus

### NEAR KANO

by R.E.Sharland.

There have been several records of Long-logged Buzzards Buteo rufinus in Nigeria in recent years (BNOS 1 (4): 15; 5:45 & 53). In February 1968, R.J.Dowsett and I found one in an area of waste land upon which are a large number of wireless masts. This bird stayed until the end of March. In each subsequent year I have observed a Long-logged Buzzard in this area and this winter (1971/72) there have been two birds.

In January 1970 I discovered another Long-legged Buzzard in an area of scrub, seventeen miles ENE of Kano, at Gesawa (12 81N., 8 451H), and here too buzzards have been observed in every subseqt ent winter.

All the specimens observed to date have been pale phased birds, with pale heads and very pale, unbarred, tails. The underwing pattern is striking with whitish flight feathers which are dark tipped and a darkish patch at the carpal joint.

When disturbed the birds usually fly slowly from one Baobab tree to another, although once (at Gesawa) I did observe a buzzard soaring.